



Kawm Ntawv Xyoo _____

KEV PAB THAUS MENYUAM PHIV (Phiv Kab, Zaubmov los Yas)

Menyuam Npe _____ Hnub Yug _____
Tsev Kawm Ntawv _____ Qib _____ Naikhu/Homeroom _____
Niamtxiv/Tus Saib Xyuas _____
Chaw Nyob _____
Xovtooj(Hauv tsev) _____ Xovtooj(Ntawm tes) _____ Xovtooj(Haujlwm) _____
Tivtauj thaum Xwmceev _____ Txheeb li cas _____
Xovtooj(Hauv tsev) _____ Xovtooj(Ntawm tes) _____ Xovtooj(Haujlwm) _____
Kws Kho Mob _____ Xovtooj _____

***Zaub mov phiv/tsi haum** lub cev zom tsi tau qee yam zaub mov, lossis yam zaub mov ntawd mus ua rau tej plab hnyuv ua haujlwm tsi zoo. Cov cim ua phiv zaub mov xwsli xeevsiab, tsam plab, mob plab, raws plab, tsi tau zoo nyob lossis mob taubhau.

Kuv tus menyuam TSI MUAJ khoom phiv, tiamsi muaj tej zaubmov noj tsi haum: _____

☐ Txwv txhua lub sijhawm ☐ pub meme rau los tau ☐ Tsi txwv hauv tsev kawm ntawv

Thaus noj zaubmov tsi haum, kuv tus menyuam cov cim mob yog: _____

***Kev Phiv Zaubmov** tshwmsim thaum cov roj ntsha tua kabmov hauv lub cev pom tias zaubmov yog yam uas los tawmtsam nws. Qhov no dhau mus ua khoom phiv thiab tshwm tej cim mob xwsli ua xua, ntuav, mob plab, cajpas nruj nruj, hais lus hawb hawb, hnoos, teebmeem ua pa, ntshav poob qis. Ua rau tuag tau.

PAB KHO KEV PHIV

Kuv tus menyuam phiv: _____

Yog koj tus menyuam phiv yas, puas tshem tej yas tawm ib ncig ntawm nws hauv tsev kawm ntawv mus? ☐ **Tshem** ☐ **Tsi Tshem**

Khij cov cim mob koj tus menyuam muaj dhau los lawm thaum nws phiv:

☐ Pob Xua ☐ Khaus ☐ Hauv Siab Ceev Ceev ☐ Ua Pa Nyuaj
☐ Nplais su/o ☐ Kiv taubhaus ☐ Ntshav Poob Qis ☐ Tsi Hnov
☐ Lub chaw ntawd su/o ☐ Lub chaw ntawd liab liab ☐ Lwm yam _____

Yog cov cim mob tshwm tuaj, yuav tshwm sai npaum cas thiab cov neeg ua haujlwm hauv tsev kawm ntawv yuavtsum tau ua dabtsi pab rau? _____

Koj tus menyuam puas muaj txog txog siav (asthma): ☐ **Muaj** (cim mob yuav tshwm yoojyim dua) ☐ **Tsi muaj**

Menyuam hnyav: _____ lbs.

Koj tus menyuam puas yuavtsum tau siv tshuaj phiv?***Yuavtsum muaj Daim Ntawv Tso Cai Siv Tshuaj***

Koj tus menyuam puas siv tus Epi Pen?

Koj tus menyuam puas tso ib tus Epi Pen rau hauv tsev kawm ntawv?

☐ Siv ☐ Tsi siv Dabtsi? _____

☐ Siv ☐ Tsi siv

☐ Tso ☐ Tsi tso Tso Twg? _____

Yog yuavtsum tau siv tshuaj hauv tsev kawm ntawv, yuavtsum tau ua daim ntawv **Tso Cai Siv Tshuaj** kom tiav

Yog koj tus menyuam yuavtsum tau noj tej yam zaubmov txawv vim nws muaj phiv, koj **yuavtsum** hais rau **Tsev Kawm Ntawv Qhov Chaws Saib Xyuas** **Zaubmov** ntawm xovtooj (920)208-4587

Niamtxiv/Tus Saib Xyuas Kos Npe _____ **Hnub tim** _____

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS

**LUNG**

Short of breath,
wheezing,
repetitive cough

**HEART**

Pale, blue,
faint, weak
pulse, dizzy

**THROAT**

Tight, hoarse,
trouble
breathing/
swallowing

**MOUTH**

Significant
swelling of the
tongue and/or lips

**SKIN**

Many hives over
body, widespread
redness

**GUT**

Repetitive
vomiting, severe
diarrhea

**OTHER**

Feeling
something bad is
about to happen,
anxiety, confusion

**OR A
COMBINATION**
of symptoms
from different
body areas.



1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell them the child is having anaphylaxis and may need epinephrine when they arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.

MILD SYMPTOMS**NOSE**

Itchy/runny
nose,
sneezing

**MOUTH**

Itchy mouth

**SKIN**

A few hives,
mild itch

**GUT**

Mild nausea/
discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE
SYSTEM AREA, GIVE EPINEPHRINE.**


**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.


**NOTE: Do not depend on
antihistamines or inhalers
(bronchodilators) to treat a
severe reaction. USE
EPINEPHRINE!**

EPIPEN® (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS

1. Remove the EpiPen Auto-Injector from the plastic carrying case.
2. Pull off the blue safety release cap.
3. Swing and firmly push orange tip against mid-outer thigh.
4. Hold for approximately 10 seconds.
5. Remove and massage the area for 10 seconds.

**AUVI-Q™ (EPINEPHRINE INJECTION, USP) DIRECTIONS**

1. Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
2. Pull off red safety guard.
3. Place black end against mid-outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.

**HOW TO USE NEFFY® (EPINEPHRINE NASAL SPRAY)**

1. Remove neffy from packaging. Pull open the packaging to remove the neffy nasal spray device.
2. Hold device as shown. Hold the device with your thumb on the bottom of the plunger and a finger on either side of the nozzle. Do not pull or push on the plunger. Do not test or prime (pre-spray). Each device has only 1 spray.
3. Insert the nozzle into a nostril until your fingers touch your nose. Keep the nozzle straight into the nose pointed toward your forehead. Do not point (angle) the nozzle to the nasal septum (wall between your 2 nostrils) or outer wall of the nose.
4. Press plunger up firmly until it snaps up and sprays liquid into the nostril. Do not sniff during or after the dose is given. If any liquid drips out of the nose, you may need to give a second dose of neffy after checking for symptoms.
5. If symptoms don't improve or worsen within 5 minutes of initial dose, administer a second dose into the same nostril with a new neffy device.



This information may be shared with the classroom teacher(s), bus driver, and other appropriate school personnel with a need to know.

foodallergy.org

DL 080525 SI